



# X-ray spectroscopy of the brightest $\gamma$ -ray nova ASASSN-18fv



Swift/XRT images of a nova 50 to 400 days post-explosion


Kirill Sokolovsky, Elias Aydi, Laura Chomiuk,  
Adam Kawash (Michigan State University),  
Koji Mukai (NASA/GSFC), Raimundo Lopes  
(Universidade Federal de Sergipe), Thomas Nelson  
(University of Pittsburgh), Brian D. Metzger,  
Elad Steinberg (Columbia University)

# Classical novae emit X-rays




- **X-ray fireball** in first hours of explosion (never observed)
- Shock waves heat plasma and accelerate particles weeks-months after explosion
- Hydrogen-burning white dwarf - “Super-Soft Source”
- When accretion restarts, the gas hitting WD surface gets shocked and heated to X-ray temperatures

# Classical novae emit X-rays




- X-ray fireball in first hours of explosion (never observed)
- **Shock waves** heat plasma and accelerate particles weeks-months after explosion
- Hydrogen-burning white dwarf - “Super-Soft Source”
- When accretion restarts, the gas hitting WD surface gets shocked and heated to X-ray temperatures

# Classical novae emit X-rays




- **X-ray fireball** in first hours of explosion (never observed)
- **Shock waves** heat plasma and accelerate particles weeks-months after explosion
- **Hydrogen-burning white dwarf** - “Super-Soft Source”
- When **accretion** restarts, the gas hitting WD surface gets shocked and heated to X-ray temperatures

# Classical novae emit X-rays




- **X-ray fireball** in first hours of explosion (never observed)
- **Shock waves** heat plasma and accelerate particles weeks-months after explosion
- **Hydrogen-burning white dwarf** - “Super-Soft Source”
- **When accretion** restarts, the gas hitting WD surface gets shocked and heated to X-ray temperatures


# X-ray observatories



- **Swift** (0.3-10 keV): fast repointing  
-> can do long-term monitoring





- **NuSTAR** (3-78 keV): exceptional sensitivity to hard X-rays





- **XMM-Newton** & **Chandra**: can do high-resolution spectroscopy with X-ray gratings

# Nova Cir 2018: Super-Soft Source





# V392 Per: accretion-powered X-rays?








# ASASSN-18fv: shocks to Super-Soft



# ASASSN-18fv: shocks to Super-Soft



# ASASSN-18fv = N Car 2018 = V906 Car




ASASSN-18fv and  $\eta$  Car nebula imaged by Joseph Brimacombe

Discovered 2018-03-20.32 UT  
by the [ASAS-SN survey](#)





# ASASSN-18fv = N Car 2018 = V906 Car




ASASSN-18fv and  $\eta$  Car nebula imaged by Joseph Brimacombe


Discovered 2018-03-20.32 UT  
by the [ASAS-SN survey](#)



# MW observations of ASASSN-18fv



# Nova Car 2018: NuSTAR spectra



**day 36:**

$$kT = 8.6 \pm 0.9 \text{ keV}$$

$$nH = 4.3 \pm 2.3 \times 10^{22} \text{ cm}^{-2}$$

$$F = 2.7 \times 10^{-12} \text{ ergs/s/cm}^2$$

**day 57:**

$$kT = 4.4 \pm 0.2 \text{ keV}$$

$$nH = 0.6 \pm 0.3 \times 10^{22} \text{ cm}^{-2}$$

$$F = 3.5 \times 10^{-12} \text{ ergs/s/cm}^2$$

CNO overabundance w.r.t.  
solar =  $210 \pm 110$

no non-thermal emission




# NuSTAR observations of novae

Five novae observed so far:

- **V745 Sco** (WD+RG) - detected ([Orio et al. 2015](#))
- **V339 Del** - not detected (Mukai et al. in prep.)
- **V5668 Sgr** - not detected (Mukai et al. in prep.)
- **V5855 Sgr** - detected while still  $\gamma$ -ray bright ([Nelson et al. 2019](#))
- **ASASSN-18fv** - detected while still  $\gamma$ -ray bright

Consistent with thermal emission in all cases

# XMM observations of ASASSN-18fv



constant\*phabs\*vphabs\*bvapec

EPIC+RGS

PHABS

$N_H$  ( $\times 10^{21}$  cm<sup>-2</sup>)  $2.4^{+0.4}_{-0.3}$

VPHABS

$N_H$  ( $\times 10^{21}$  cm<sup>-2</sup>)  $0.12^{+0.03}_{-0.03}$

BVAPEC

$kT$  (keV)  $1.07^{+0.04}_{-0.01}$

redshift  $-2.9 \times 10^{-3*}$

velocity (km s<sup>-1</sup>)  $378^{(*)}$

$N/N_{\odot}$   $345^{+93}_{-70}$

$O/O_{\odot}$   $29^{+7}_{-5}$

$Ne/Ne_{\odot}$   $2.2^{+0.6}_{-0.5}$

$Mg/Mg_{\odot}$   $0.6^{+0.2}_{-0.1}$

$Si/Si_{\odot}$   $1.1^{+0.2}_{-0.2}$


$Fe/Fe_{\odot}$   $<0.1$

$\chi^2_{\nu}$  1.15

d.o.f. 1837



# XMM observations of ASASSN-18fv



constant\*phabs\*vphabs\*bvapec

EPIC+RGS

PHABS

$N_H$ ( $\times 10^{21} \text{ cm}^{-2}$ )	$2.4^{+0.4}_{-0.3}$
--	---------------------

VPHABS

$N_H$ ( $\times 10^{21} \text{ cm}^{-2}$ )	$0.12^{+0.03}_{-0.03}$
--	------------------------

BVAPEC

$kT$ (keV)	$1.07^{+0.04}_{-0.01}$
------------	------------------------

redshift	$-2.9 \times 10^{-3*}$
----------	------------------------

velocity ( $\text{km s}^{-1}$ )	$378^{(*)}$
---------------------------------	-------------

$N/N_{\odot}$	$345^{+93}_{-70}$
---------------	-------------------

$O/O_{\odot}$	$29^{+7}_{-5}$
---------------	----------------

$Ne/Ne_{\odot}$	$2.2^{+0.6}_{-0.5}$
-----------------	---------------------

$Mg/Mg_{\odot}$	$0.6^{+0.2}_{-0.1}$
-----------------	---------------------

$Si/Si_{\odot}$	$1.1^{+0.2}_{-0.2}$
-----------------	---------------------

$Fe/Fe_{\odot}$	$<0.1$
-----------------	--------

$\chi^2_{\nu}$	1.15
----------------	------

d.o.f.	1837
--------	------

# Outlook

- X-rays provide a **new window on nova physics**
- **Shock energy, abundances, ejecta mass, non-thermal emission**
- Constrains on **particle acceleration** from X-ray/GeV
- **Where, when** and **why shocks** are formed?
- Relevant for distant **shock-powered transients**:  
Type IIn SNe, TDEs, stellar mergers

White paper on shocks by [Chomiuk et al. \(2019\)](#)

# NuSTAR observations of ASASSN-18fv

*NuSTAR* observing log

ObsID	Epoch (days)	Start UT	Stop UT	Exposure FPMA (ks)	Exposure FPMB (ks)	Net count rate FPMA (cts/s)	Net count rate FPMB (cts/s)
80301306002	36.3	2018-04-20 14:46	2018-04-22 02:01	48.8	48.5	$0.01582 \pm 0.00066$	$0.01630 \pm 0.00067$
90401322002	57.2	2018-05-11 16:26	2018-05-12 18:01	47.5	47.4	$0.04343 \pm 0.00102$	$0.04184 \pm 0.00101$

**Column designation:** Col. 1 – observation identification number; Col. 2 – time since outburst; Col. 3 and 4 – start and stop time of the observation (interrupted by Earth occultations and South Atlantic Anomaly passes); Col. 5 and 6 – total on-source exposure time for FPMA and FPMB, respectively; Col. 7 and 8 – source count rate (background-subtracted) for FPMA and FPMB, respectively.

constant\*vphabs\*vaptec model for the two *NuSTAR* observations

Epoch (days)	$n_{HI}$ ( $\times 10^{22} \text{ cm}^{-2}$ )	$kT$ (keV)	CNO abundances	$C_{\text{FPMB}}$	Model 3.5-78.0 keV flux $\log_{10}(\text{ergs/cm}^2/\text{s})$
$\chi^2_{\text{red}} = 1.0457$ , d.o.f. = 199, $p = 0.31$					
36	$4.287 \pm 2.288$	$8.59 \pm 0.88$	$209.6 \pm 110.4$	$1.107 \pm 0.062$	$-11.564 \pm 0.012$
57	$0.568 \pm 0.288$	$4.38 \pm 0.17$		$1.006 \pm 0.034$	$-11.454 \pm 0.007$

	Case 1	Case 2	Case 3	Case 4	Case 5
	EPIC+RGS	EPIC+RGS	EPIC+RGS	RGS	RGS
PHABS					
$N_H$ ( $\times 10^{21}$ cm $^{-2}$ )	$1.8^{+0.3}_{-0.2}$	$1.8^{+0.2}_{-0.2}$	$2.4^{+0.4}_{-0.3}$	$2.1^{+0.5}_{-1.0}$	$2.0^{+2.1}_{-1.0}$
VPHABS					
$N_H$ ( $\times 10^{21}$ cm $^{-2}$ )	$0.08^{+0.02}_{-0.02}$	$0.13^{+0.03}_{-0.02}$	$0.12^{+0.03}_{-0.03}$	<0.4	<0.4
BVAPEC					
$kT$ (keV)	$1.06^{+0.01}_{-0.01}$	$1.11^{+0.01}_{-0.01}$	$1.07^{+0.04}_{-0.01}$	$0.79^{+0.04}_{-0.10}$	$0.98^{+0.15}_{-0.12}$
redshift	$(-2.9 \pm 0.1) \times 10^{-3}$	$(-2.9 \pm 0.2) \times 10^{-3}$	$-2.9 \times 10^{-3*}$	$(-3.1 \pm 0.2) \times 10^{-3}$	$-2.9 \times 10^{-3(*)}$
velocity (km s $^{-1}$ )	$394 \pm 70$	$378 \pm 72$	$378^{(*)}$	$386^{+72}_{-76}$	$378^*$
N/N $_{\odot}$	$728^{+232}_{-150}$	$403^{+99}_{-73}$	$345^{+93}_{-70}$	$230^{+236}_{-81}$	$212^{+197}_{-87}$
O/O $_{\odot}$	$30^{+7}_{-6}$	$24^{+4}_{-5}$	$29^{+7}_{-5}$	$14^{+15}_{-5}$	$17^{+12}_{-5}$
Ne/Ne $_{\odot}$	$0.7^{+0.6}_{-0.5}$	$2.3^{+0.6}_{-0.5}$	$2.2^{+0.6}_{-0.5}$	$1.1^{+1.3}_{-0.5}$	$1.5^{+1.3}_{-0.7}$
Mg/Mg $_{\odot}$	$1.0^{+0.2}_{-0.2}$	$0.7^{+0.2}_{-0.1}$	$0.6^{+0.2}_{-0.1}$	$1.0^{+1.0}_{-0.3}$	$0.9^{+0.6}_{-0.3}$
Si/Si $_{\odot}$	$1.6^{+0.4}_{-0.3}$	$1.2^{+0.2}_{-0.2}$	$1.1^{+0.2}_{-0.2}$	$1.0^{+2.1}_{-0.7}$	$2.0^{+1.3}_{-0.5}$
Fe/Fe $_{\odot}$	$0.17^{+0.08}_{-0.05}$	<0.1	<0.1	<0.13	<0.04
$\chi^2_{\nu}$	1.25	1.16	1.15	1.01	1.01
d.o.f.	1847	1488	1837	987	977

Notes:

Model CONSTANT\*PHABS\*VPHABS\*BVAPEC in four different cases:

Case 1: in the whole spectral coverage, without Gaussian lines;

Case 2: excluding spectral regions associated with (r,i,f) lines: 0.4-0.45 keV, 0.55-0.6 keV, 0.85-0.95 keV, and 1.3-1.4 keV;


Case 3: in the whole spectral coverage, including Gaussian lines associated with r,i,f lines (Table 3);

Case 4: only RGS, without Gaussian lines;

Case 5: only RGS, with Gaussian lines associated with r,i,f lines (Table 3);

Abundance table: aspl: Asplund M, Grevesse N., Sauval A.J. & Scott P., 2009, ARAA, 47, 481;

~100 observed,  
60% detected,  
a few lightcurves

Ness et al. (2007),  
Schwarz et al. (2011),   
Osborne (2015)

**hard -> med. -> soft**



2011).